

國立清華大學 100 學年度碩士班入學考試試題

系所班組別：工程與系統科學系乙組

考試科目（代碼）：工程數學(2901)


共 3 頁，第 1 頁 \*請在【答案卷、卡】作答

1. Find the transient current if

$$R = 6 \Omega, L = 1 H, C = 0.04 F, E = 600 (\cos t + 4 \sin t) V;$$

(L, R, C, E, are measured in henrys, ohms, farads, volts, respectively.)

Initial current and charge are assumed to be zero. (10%)



2. Consider a general nonhomogeneous linear ODE

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + p_{n-2}(x)y^{(n-2)} + \dots + p_1(x)y' + p_0(x)y = r(x)$$

The particular solution  $y_p(x)$  can be solved by the method of variation of parameters. That is,

$$y_p(x) = \sum_{k=1}^n y_k(x) \int \frac{W_k(x)}{W(x)} r(x) dx$$

(Where the  $y_k$ 's are  $n$  linearly independent homogeneous solutions.  $W$  is the Wronskian of  $y_1, \dots, y_n$ , and  $W_k$  is identical to  $W$ , but with the  $k$ th column replaced by a column of zeros-except for the bottom element, which is 1.)

Try to solve the following equation using the method of variation of parameters.

$$y''' + \frac{3}{4}x^{-2}y' - \frac{3}{4}x^{-3}y = 9x^{5/2} \quad (10\%)$$

國立清華大學 100 學年度碩士班入學考試試題

系所班組別：工程與系統科學系乙組

考試科目（代碼）：工程數學(2901)

共 3 頁，第 2 頁 \*請在【答案卷、卡】作答

3. Solve the following equation by Laplace Transform method.

$$y'+y = f(t), \quad y(0) = 3,$$
$$\text{where } f(t) = \begin{cases} 0 & 0 \leq t < \pi \\ 2 \cos t & t \geq \pi \end{cases}. \quad (10\%)$$

4. Find the basis of solutions  $y(x)$  of the following differential equation. Show the details of your work.

$$xy'' + (2x+1)y' + (x+1)y = 0. \quad (10\%)$$

5. Find a unit vector normal to surface  $S$  given by  $\cos(xy) = e^z - 1$  at the point  $(1, \pi, 0)$ . (10%)

6. Let  $\mathbf{F} = (x-y)\mathbf{i} + (y-z)\mathbf{j} + (z-x)\mathbf{k}$ . Evaluate the surface integral of  $\mathbf{F}$  over the unit sphere defined by  $x^2 + y^2 + z^2 = 1$ . (10%)

7. Define the Fourier transform of  $f(x)$  to be  $\hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{-iwx} dx$ .

(a) [5%] Calculate the Fourier transform of  $f(x) = \begin{cases} a-|x| & , |x| < a \\ 0 & , \text{otherwise} \end{cases}$

(b) [5%] Consider the one-dimensional diffusion equation:

$$\frac{\partial}{\partial t} u(x,t) = D \frac{\partial^2}{\partial x^2} u(x,t) \quad \text{for } -\infty < x < \infty$$

with the initial condition  $u(x,0) = f(x)$ . Use the Fourier transform to show that the

solution of the diffusion equation takes the form  $u(x,t) = \int_{-\infty}^{\infty} K(x-\xi,t) f(\xi) d\xi$ .

Find  $K(x-\xi,t)$ , which is called the kernel.

[Hint: Gaussian integral  $\int_{-\infty}^{\infty} \exp\left(\frac{-x^2}{2\sigma^2}\right) dx = \sqrt{2\pi\sigma^2}$  ]

國立清華大學 100 學年度碩士班入學考試試題

系所班組別：工程與系統科學系 乙組

考試科目（代碼）：工程數學(2901)

共 3 頁，第 3 頁 \*請在【答案卷、卡】作答

8. (a) [5%] Find and classify all local maxima, local minima and saddles for

$$f(x, y, z) = \exp(2x^2 + xz - 5z^2).$$

(b) [5%] Consider a forced vibration system which is described by the equations

$$\begin{aligned} \frac{d^2 x_1}{dt^2} + 2x_1 - x_2 &= A \sin(\omega t) \\ \frac{d^2 x_2}{dt^2} - x_1 + 2x_2 &= B \sin(\omega t) \end{aligned}, \text{ where } A, B, \text{ and } \omega \text{ are constant.}$$

To seek a particular solution, we assume  $x_1(t) = q_1 \sin(\omega t)$  and  $x_2(t) = q_2 \sin(\omega t)$ .

Find  $q_1$  and  $q_2$ .

9. Along the circumference of the circle  $r = b$  a solution  $T(r, \theta)$  of Laplace's equation is required to take on the value  $T_0$  when  $0 < \theta < \pi$  and the value  $-T_0$  when  $\pi < \theta < 2\pi$ . Determine an expression for  $T$  valid when  $r > b$ . (Show the details of your work.)

$$[\nabla^2 T = \frac{1}{r} \frac{\partial}{\partial r} (r \frac{\partial T}{\partial r}) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \theta^2}] \quad (12\%)$$

10. (complex analysis) Prove the following identity

$$\sin^{-1} z + \cos^{-1} z = \frac{1}{2}(4n+1)\pi, \quad n = 0, \pm 1, \pm 2, \dots \quad (8\%)$$